

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

BRAVECTO[®]

112.5 MG FLURALANER Chewable Tablets for Very Small Dogs
250 MG FLURALANER Chewable Tablets for Small Dogs
500 MG FLURALANER Chewable Tablets for Medium Dogs
1000 MG FLURALANER Chewable Tablets for Large Dogs
1400 MG FLURALANER Chewable Tablets for Very Large Dogs

ACTIVE CONSTITUENT: 136.4 g/kg FLURALANER

BRAVECTO Chewable Tablets provide

- Treatment and prevention of flea (*Ctenocephalides felis*) infestations for **3 months**.
- Treatment and control of paralysis tick (*Ixodes holocyclus*) for **4 months**.
- Treatment and control of brown dog tick (*Rhipicephalus sanguineus*) for **2 months**.
- Treatment and control of bush tick (*Haemaphysalis longicornis*) for **2 months**.
- Rapid onset of action kills newly emerged adult fleas before they lay eggs.
- Control of flea allergy dermatitis (FAD).
- Prevention of flea tapeworm infestations.
- Highly palatable so easy to administer.

BRAVECTO Chewable Tablet is effective for the treatment and control of ear mite (*Otodectes cynotis*), demodex mite (*Demodex* spp.) and sarcoptic mange (*Sarcoptes scabiei*) after a single treatment.

BRAVECTO contains fluralaner, a member of the antiparasitic class of isoxazoline-substituted benzamide derivatives. Fluralaner acts antagonistically on arthropods ligand-gated chloride channels (GABA-receptor and glutamate-receptor). **BRAVECTO** is effective against fipronil-resistant strains of fleas.

BRAVECTO can be used in breeding, pregnant and lactating dogs.

ONSET OF ACTION AND EFFICACY

BRAVECTO provides effective control of pre-existing infestations of paralysis ticks (*Ixodes holocyclus*) within 24 hours.

BRAVECTO provides effective control of fleas on dogs within 8 hours of administration, and subsequently maintains persistent efficacy against flea re-infestations for **3 months** following administration. The flea life cycle can be broken by the rapid onset of action and the long-lasting efficacy against adult fleas on the dog. Newly emerged adult fleas on a dog are killed before viable eggs are produced. Effective flea control with **BRAVECTO** Chewable Tablets prevents flea tapeworm (*Dipylidium caninum*) infestation in dogs. A single treatment of **BRAVECTO** Chewable Tablet prevents flea tapeworm (*Dipylidium caninum*) transmission from infected fleas to susceptible dogs for up to 3 months.

BRAVECTO can be used as part of a treatment strategy for flea allergy dermatitis (FAD). *Ixodes holocyclus* ticks do not occur in Western Australia.

In tick season daily searching for and removal of any ticks found is recommended.

DIRECTIONS FOR USE

Precautions

BRAVECTO should not be used in puppies less than 8 weeks of age or weighing less than 2 kg. Use with caution in dogs with pre-existing epilepsy.

Dosage and administration

BRAVECTO is available in five sizes. The following table indicates the size of chewable tablet to be used according to the body weight of the dog (equivalent to 25-56 mg fluralaner/kg body weight).

Body weight of dog (kg)	Chewable tablet size to be used
2 - 4.5	BRAVECTO 112.5 MG FLURALANER Chewable Tablets for Very Small Dogs
>4.5 - 10	BRAVECTO 250 MG FLURALANER Chewable Tablets for Small Dogs
>10 - 20	BRAVECTO 500 MG FLURALANER Chewable Tablets for Medium Dogs
>20 - 40	BRAVECTO 1000 MG FLURALANER Chewable Tablets for Large Dogs
>40 - 56	BRAVECTO 1400 MG FLURALANER Chewable Tablets for Very Large Dogs

Within each weight range, a whole tablet must be used. For dogs weighing more than 56 kg, use a combination of two tablets that most closely matches the body weight.

In dogs older than 8 weeks, treatment every 2 months has been shown to be well tolerated.

Puppies should be weighed regularly. Fast growing puppies that outgrow the initial weight band during the retreatment interval, may be retreated for fleas and paralysis ticks at 2 monthly intervals. Treatment may be appropriately adapted by the veterinarian to suit the individual weight changes.

Administer **BRAVECTO** at or around the time of feeding. **BRAVECTO** is readily consumed by most dogs when given by the owner. **BRAVECTO** can also be given with food or administered in the same way as other tablets. **BRAVECTO** chewable tablets can be used year round.

Treatment schedule

For optimal treatment and control of **flea infestation**, use **BRAVECTO** Chewable Tablet every **3 months**.

For optimal treatment and control of **paralysis tick**, use **BRAVECTO** Chewable Tablet every **4 months**.

For optimal treatment and control of **brown dog tick** and **bush tick**, use **BRAVECTO** Chewable Tablet every **2 months**.

One dose of **BRAVECTO** Chewable Tablet clears **ear mite** and **sarcoptic mange** infestations within **1 month**, and **demodex mite** infestation within **2 months**. The absence of mites can be confirmed by two consecutive monthly skin scrapings. If mite infestation reoccurs, consult with a veterinarian.

General directions

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

Fluralaner is part of the isoxazoline family of chemicals. Adverse reactions to this family of chemicals are rarely observed but may include vomiting, diarrhoea, lethargy, inappetence, itching and very rarely, seizures. Most adverse reactions are self-limiting and of short duration. If you have any concerns, please speak to your veterinarian.

Concurrent treatment of **BRAVECTO** and milbemycin oxime / praziquantel tablets is well tolerated.

Clinical studies have demonstrated that **BRAVECTO** is well tolerated in dogs when co-administered with macrocyclic lactones, and is also well tolerated in ivermectin-sensitive Collies.

For paralysis tick control it is recommended that the product be used in conjunction with daily searching for, and removal of, any ticks found as the risk of tick paralysis cannot be entirely eliminated. Speak with your veterinarian for advice on tick searching and removal techniques.

For further information contact MSD Animal Health Customer Service on 1800 230 833.

SAFETY DIRECTIONS

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. *Phone Australia 131126.*

DISPOSAL

Dispose of empty foil and carton by wrapping in paper and putting in garbage.

STORAGE

Store below 30°C (room temperature).

APVMA

BRAVECTO 112.5 MG FLURALANER Chewable Tablets for Very Small Dogs - 68867/119011
BRAVECTO 250 MG FLURALANER Chewable Tablets for Small Dogs - 68872/119010
BRAVECTO 500 MG FLURALANER Chewable Tablets for Medium Dogs - 68871/119009
BRAVECTO 1000 MG FLURALANER Chewable Tablets for Large Dogs - 68870/119008
BRAVECTO 1400 MG FLURALANER Chewable Tablets for Very Large Dogs - 68873/119006

Warranty

Intervet Australia Pty Limited (IAPL), trading as MSD Animal Health, warrants that this product is of merchantable quality and fit for its intended purpose. IAPL's liability for any loss, including consequential losses or injury caused by any act or omission, including negligent acts or omissions, by IAPL or its agent, is limited to replacing or repairing the product at the option of IAPL. If possible, a sample of any product causing concern should be retained or delivered to IAPL within 30 days for a scientific examination.

Intervet Australia Pty Limited
(trading as MSD Animal Health)
91-105 Harpin Street,
Bendigo East VIC 3550
Phone: 1800 230 833
www.bravecto.com.au

© Registered trademark

© Copyright 2019

341964 R2



