

# MICROLAX ENEMA

## PRODUCT SAFE USE INFORMATION SHEET.

DATASHEET BY – <http://www.familybrands.co.nz>

Gentle relief of constipation usually within 30 minutes.

### CLASSIFICATION

Pharmacy Medicine

### INGREDIENTS

Microenema 5 mL

### HOW TO USE

1. Twist and pull the seal off the nozzle. 2. Squeeze the tube slightly so that a drop of Microlax smears the tip and thus makes insertion easier.
3. Insert the nozzle fully into the rectum. N.B. In children under 3 years of age insert only half the length of the nozzle (see indication on the nozzle).
4. Squeeze out the contents fully by squeezing the shoulder of the tube.
5. Keeping the tube squeezed tightly, withdraw the nozzle.

### WARNINGS AND PRECAUTIONS

Microlax should be kept in a place where the temperature stays below 30°C.  
Do not store Microlax or any other medicine in the bathroom or near a sink.  
KEEP MICROLAX OUT OF REACH OF CHILDREN.

### SIDE EFFECTS

Microlax, like all medicines, may cause unwanted side effects in some people.  
A slight rectal burning sensation has occasionally been reported. Individual cases of allergy or abnormal reaction (general skin reaction with or without low blood pressure or breathing difficulties) have also been reported.  
If these effects do not go away or they are worrying you, tell your doctor or pharmacist.  
Some people may get other side effects while using Microlax.